arXiv:0906.1035v2 [math.DG] 24 Mar 2010

Ricci Flow on 3-dimensional Lie groups and 4-dimensional Ricci-flat manifolds

Kensuke Onda*

Graduate School of Mathematics, Nagoya University

Abstract

We study relation of the Ricci Flow on 3-dimensional Lie groups and 4-dimensional Ricci-flat manifolds. In particular, we construct Ricci-flat cohomogeneity one metrics for 3 dimensional Lie groups.

1 Introduction

In this paper, we will discuss some relationship between the Ricci flow of left-invariant Riemannian metrics on 3-dimensional unimodular simply-connected Lie group G and the Ricci-flat metrics of cohomogeneity one on the space-time $\mathbb{R} \times G$.

First of all, we introduce the notion of cohomogeneity one metrics with respect to a Lie group G.

Definition 1.1. A pseudo-Riemannian manifold (M, g) is a cohomogeneity one with respect to a Lie group G, if and only if G is a subgroup of $\mathrm{Isom}(M, g)$, and the codimension of principal orbits under the action of G equals 1.

The simplest example of a cohomogeneity one metric arises from the standard action of $SO(n)$ to $\mathbb{R}^n$. The singular orbit is $\{0\}$, and the principal orbits are S^{n-1} of various radii.

---

*kensuke.onda@math.nagoya-u.ac.jp

In this paper, we attempt to construct cohomogeneity one Einstein metrics (in particular Ricci-flat metrics) from the Ricci flow solutions of left-invariant metrics on 3-dimensional unimodular simply connected Lie groups. The Ricci-flat metrics obtained in this paper has the property that their sectional curvatures decay to 0 at one end toward which the metric is complete. Such metrics was studied by Lorentz, Gibbons, Hawking, Pope and many people. Some of them is called ALF and the Taub-NUT metric on $\mathbb{R} \times SU(2)$ arising from the Ricci flow of the left $SU(2)$- and right $U(1)$-invariant metrics on $SU(2)$ is ALF.

Before proceeding to general 3-dimensional Lie groups, we examine the case of $SU(2)$, which is well known. There exists a left-invariant coframe $\{\theta^i\}_{i=1}^3$ on $SU(2)$ satisfying $d\theta^i = 2\theta^j \wedge \theta^k$, where (i,j,k) are cyclic permutation of $\{1,2,3\}$. Then cohomogeneity one metrics with respect to $SU(2)$ is described as

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{1.1}$$

Some special cases of (1.1) are listed in the following examples:

1. If $a = b = c$ are linear, then the metric g has constant curvature 0.
2. If a, b and c are constants, then the metric g becomes a product metric on $\mathbb{R} \times SU(2)$.
3. If $a = b = c = \sin t$, then the metric g has positive constant curvature.
4. If $a = b = c = \sinh t$, then the metric g has negative constant curvature.

Thus, it is interesting to ask for which $\{a(t), b(t), c(t)\}$ the resulting cohomogeneity one metric is Einstein. In this paper, when the triple of functions $\{a(t), b(t), c(t)\}$ satisfies the Ricci flow equation, we examine whether the resulting cohomogeneity one metric is a Ricci-flat metric. This generalizes the above example 1.

First of all, we define the *Ricci flow* again. In this paper, we define that a 1-parameter family $g(t)$ of Riemannian metrics is the Ricci flow if and only if it solves

$$\frac{\partial}{\partial t} g(t)_{ij} = -\mathrm{Ric}[g(t)]_{ij},$$

and a 1-parameter family of Riemannian metrics $g(t)$ is the backward Ricci flow if and only if it solves

$$\frac{\partial}{\partial t} g(t)_{ij} = \mathrm{Ric}[g(t)]_{ij}.$$

In the Introduction, the Ricci flow was defined by

$$\frac{\partial}{\partial t} g(t)_{ij} = -2\mathrm{Ric}[g]_{ij}.$$

The difference in 1 and 2 is just a choice of the scale and produces no trouble. Indeed, if we put $h := \frac{k}{2} g$, where k is a positive constant, then

$$\frac{\partial}{\partial t} h(t)_{ij} = (\frac{k}{2})(-2\mathrm{Ric}[g]_{ij}) = -k\mathrm{Ric}[h]_{ij},$$

because $\mathrm{Ric}[h] = \mathrm{Ric}[\frac{k}{2} g] = \mathrm{Ric}[g]$. In this paper, if k equals 1, we say that $h(t)$ is a solution to the Ricci flow. Also the backward Ricci flow (in this paper) is defined similarly. Note that if we change t into $-t$, the Ricci flow equation changes into the backward Ricci flow equation.

Next, we review the Ricci flow on $SU(2)$. Let $\{F_i\}_{i=1}^3$ be an orthonormal frame of a left-invariant metric g_3 on $SU(2)$, satisfying $[F_i, F_j] = -2F_k$ $(i, j, k$: cyclic), and $\{\theta^i\}$ the dual coframe of $\{F_i\}_{i=1}^3$. The left-invariant metric g_3 is expressed as

$$g_3 = A(\theta^1)^2 + B(\theta^2)^2 + C(\theta^3)^2.$$

Then the Ricci flow is equivalent to the system of ODE's:

$$\begin{aligned} \frac{d}{dt} A &= \frac{(B-C)^2 - A^2}{BC}, \\ \frac{d}{dt} B &= \frac{(C-A)^2 - B^2}{CA}, \\ \frac{d}{dt} C &= \frac{(A-B)^2 - C^2}{AB}. \end{aligned} \tag{1.2}$$

The behavior of solutions of (1.2) is known in [KM01, CK04].

Proposition 1.2 ([KM01, CK04]). *The solution of the Ricci flow equation exists on $(-\infty, T)$, where T depends on the initial data, and if t goes to T, then g_3 becomes asymptotically round and shrinks to a point.*

Next we consider a cohomogeneity one metric with respect to $SU(2)$, given by

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{1.3}$$

A typical example of cohomogeneity one metric with respect to $SU(2)$ is the Taub-NUT metric, given by

$$g = \Big(\frac{r+m}{r-m}\Big)dr^2 + (r^2-m^2)\{(\theta^1)^2+(\theta^2)^2\} + 4m^2\Big(\frac{r-m}{r+m}\Big)(\theta^3)^2.$$

As is well known, this metric has the following properties. First, this metric is a hyper-Kähler metric, and therefore this is a Ricci-flat metric. Secondly, we can put

$$a = b = (r^2-m^2)^{\frac{1}{2}}, \quad c = 2m\Big(\frac{r-m}{r+m}\Big)^{\frac{1}{2}},$$

and therefore the coefficient a equals b, but a is not equal to c (i.e., the metric on $SU(2)$ part is left $SU(2)$- and right $U(1)$-invariant). Thirdly, the change of variables $h = \Big(\frac{r+m}{r-m}\Big)^{\frac{1}{2}}$ and $dt = hdr$ implies that this metric is regarded as a cohomogeneity one metric. Fourthly, it is easy to check that a, b and c satisfy

$$\begin{aligned} \frac{d}{dt}a &= \frac{a^2-(b-c)^2}{bc}, \\ \frac{d}{dt}b &= \frac{b^2-(c-a)^2}{ca}, \\ \frac{d}{dt}c &= \frac{c^2-(a-b)^2}{ab}. \end{aligned}$$

This system is equivalent to the backward Ricci flow equation for left-invariant metrics on $SU(2)$. Thus we conclude that if coefficients $\{a(t), b(t), c(t)\}$ move along the Ricci flow or the backward Ricci flow on $SU(2)$, then the resulting cohomogeneity one metric (1.3) is the Taub-NUT metric and therefore Ricci-flat.

Theorem 1.3 ([Fri85, CGLP04])**.** *If a, b and c satisfy the Ricci flow equations or the backward Ricci flow equations of a metric*

$$g_3 = a(\theta^1)^2 + b(\theta^2)^2 + c(\theta^3)^2 \tag{1.4}$$

on $SU(2)$, then the cohomogeneity one metric

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2 \tag{1.5}$$

with respect to $SU(2)$ on the space-time of the Ricci flow becomes a Ricci-flat metric.

It is important to pay attention to coefficients. Coefficients of metric (1.4) are a, b and c. But, coefficients of metric (1.5) are a^2, b^2 and c^2. Therefore, a, b and c are positive (because we consider left-invariant Riemannian metrics on $SU(2)$). Even if a, b and c are positive, we have still freedom in introducing minus sign before $\{a^2, b^2, c^2\}$ in the attempt to discover cohomogeneity one metrics on the space-time of the Ricci flow (this freedom is really essential in the case of $E(1,1)$ and $SL(2,\mathbb{R})$).

Remark 1.4. Even if a cohomogeneity one metric

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2$$

with respect to $SU(2)$ is Ricci-flat, it may not satisfy the Ricci flow equation nor the backward Ricci flow equation. We consider the Eguchi-Hanson metric as a typical example. The Eguchi-Hanson metric is given by

$$g = \frac{dr^2}{1-(m/r)^4} + r^2\{(\theta^1)^2 + (\theta^2)^2\} + r^2(1-(m/r)^4)(\theta^3)^2. \tag{1.6}$$

Put

$$a = b = r,\ c = r\Big(1 - (\frac{m}{r})^4\Big)^{\frac{1}{2}}.$$

By the coordinate transform,

$$h = \frac{1}{\Big(1-(\frac{m}{r})^4\Big)^{\frac{1}{2}}},\ dt = hdr,$$

the metric (1.6) is regarded as a cohomogeneity one metric. Also this metric (1.6) is a Ricci-flat metric, but coefficients a, b and c satisfy the following system of ODE's:

$$\begin{aligned}\frac{d}{dt}a &= \frac{a^2-(b-c)^2}{bc} + 2,\\ \frac{d}{dt}b &= \frac{b^2-(c-a)^2}{ca} + 2,\\ \frac{d}{dt}c &= \frac{c^2-(a-b)^2}{ab} + 2.\end{aligned}$$

These equations are neither the Ricci flow nor the backward Ricci flow.

We now study cohomogeneity one metrics constructed from the Ricci flow solution on other groups, and construct Ricci-flat metrics on their space-time. Ricci-flat metrics that we construct in this paper are listed in Table 1.

Lie groups	the Ricci flow equation	signature of metric
$SU(2)$	$SU(2)$	$(4,0)$
$E(2)$	$E(2)$	$(4,0)$
$SL(2,\mathbb{R})$	$SU(2)$	$(2,2)$
H_3	H_3	$(4,0)$
$E(1,1)$	$E(1,1)$	$(2,2)$

Table 1: Ricci-flat metrics of cohomogeneity one corresponding to 3-dimensional Lie groups

2 Preparation

We present the definition of a Milnor frame.

Definition 2.1. Let $\{F_i\}_{i=1}^3$ be a left-invariant moving frame on G. If $\{F_i\}_{i=1}^3$ satisfies

$$[F_2,F_3]=n_1F_1,\quad [F_3,F_1]=n_2F_2,\quad [F_1,F_2]=n_3F_3,$$

where $n_i \in \{\pm 1, 0\}$, then $\{F_i\}_{i=1}^3$ is called *a Milnor frame* .

As is well known, 3-dimensional unimodular simply-connected Lie groups were classified by Milnor [Mil76].

Proposition 2.2 ([Mil76]). *Let $\{F_i\}$ be a Milnor frame. For signatures of $\{n_i\}$, 3-dimensional unimodular simply-connected Lie groups are determined as Table 2.*

Signature	Lie groups	description
$(-1,-1,-1)$, $(+1,+1,+1)$	$SU(2)$	simple
$(-1,-1,+1)$, $(-1,+1,+1)$	$\widetilde{\mathrm{SL}(2,\mathbb{R})}$	simple
$(-1,-1,0)$, $(+1,+1,0)$	$E(2)$	solvable
$(-1,0,+1)$	$E(1,1)$	solvable
$(-1,0,0)$, $(+1,0,0)$	H_3	nilpotent
$(0,0,0)$	$\mathbb{R}\oplus\mathbb{R}\oplus\mathbb{R}$	commutative

Table 2: 3-dimensional unimodular simply-connected Lie groups

Remark 2.3. We may change order of n_i.

Let (M^4, g) be a cohomogeneity one 4-dimensional manifold with respect to 3-dimensional Lie group G. Let $\{F_i\}_{i=1}^3$ be a Milnor frame of G, and $\{\theta_i\}$ the dual coframe of $\{F_i\}$. Then the metric is expressed as follows:

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{2.1}$$

From Theorem 1.1 (Koszul's formula), the Levi-Civita connection is expressed as below.

Proposition 2.4. *Let g be the cohomogeneity one metric (2.1). Then the Levi-Civita connection is given by*

$$\begin{pmatrix} 0 & \frac{\dot a}{a}F_1 & \frac{\dot b}{b}F_2 & \frac{\dot c}{c}F_3 \\ \frac{\dot a}{a}F_1 & -a\dot a F_0 & \frac{1}{2}\cdot\frac{-n_1a^2+n_2b^2+n_3c^2}{c^2}F_3 & \frac{1}{2}\cdot\frac{-n_3c^2+n_1a^2-n_2b^2}{b^2}F_2 \\ \frac{\dot b}{b}F_2 & \frac{1}{2}\cdot\frac{-n_1a^2+n_2b^2-n_3c^2}{c^2}F_3 & -b\dot b F_0 & \frac{1}{2}\cdot\frac{-n_2b^2+n_3c^2+n_1a^2}{a^2}F_1 \\ \frac{\dot c}{c}F_3 & \frac{1}{2}\cdot\frac{-n_3c^2+n_1a^2+n_2b^2}{b^2}F_2 & \frac{1}{2}\cdot\frac{-n_2b^2+n_3c^2-n_1a^2}{a^2}F_1 & -c\dot c F_0 \end{pmatrix} \tag{2.2}$$

Hence we obtain the Ricci tensor;

$$\mathrm{Ric}(F_0, F_0) = R_{00} = -\frac{\ddot a}{a} - \frac{\ddot b}{b} - \frac{\ddot c}{c}, \tag{2.3a}$$

$$\mathrm{Ric}(F_1, F_1) = R_{11} = -a\frac{(\dot a bc)^{\cdot}}{bc} - \frac{(n_2b^2 - n_3c^2)^2 - n_1^2a^4}{2b^2c^2}, \tag{2.3b}$$

$$\mathrm{Ric}(F_2, F_2) = R_{22} = -b\frac{(\dot b ca)^{\cdot}}{ca} - \frac{(n_3c^2 - n_1a^2)^2 - n_2^2b^4}{2c^2a^2}, \tag{2.3c}$$

$$\mathrm{Ric}(F_3, F_3) = R_{33} = -c\frac{(\dot c ab)^{\cdot}}{ab} - \frac{(n_1a^2 - n_2b^2)^2 - n_3^2c^4}{2a^2b^2}, \tag{2.3d}$$

and other components are 0. Remark that if a, b and c are constants, then the Ricci tensor R_{11}, R_{22} and R_{33} of the metric (2.1) are the Ricci tensors of a metric

$$g = a^2(\theta^1)^2 + b^2(\theta^2)^2 + c^2(\theta^3)^2.$$

3 The Heisenberg group

In this section, we consider the Heisenberg group H_3 and a cohomogeneity one with respect to H_3. Let $\{F_i\}$ be the Milnor frame of H_3, satisfying

$[F_1, F_2] = 0$, $[F_2, F_3] = F_1$ and $[F_3, F_1] = 0$; and $\{\theta_i\}$ the dual coframe of $\{F_i\}$. Then a left-invariant metric is expressed as

$$g = a(\theta^1)^2 + b(\theta^2)^2 + c(\theta^3)^2. \tag{3.1}$$

Therefore the Ricci flow equation is given by

$$\frac{d}{dt}a = -\frac{a^2}{2bc}, \tag{3.2a}$$
$$\frac{d}{dt}b = \frac{a}{2c}, \tag{3.2b}$$
$$\frac{d}{dt}c = \frac{a}{2b}. \tag{3.2c}$$

These equations were solved in [KM01] as follows:

$$\begin{cases} a(t) = a_0^{\frac{2}{3}} b_0^{\frac{1}{3}} c_0^{\frac{1}{3}} (\frac{3}{2}t + b_0c_0/a_0)^{-\frac{1}{3}}, \\ b(t) = a_0^{\frac{1}{3}} b_0^{\frac{2}{3}} c_0^{-\frac{1}{3}} (\frac{3}{2}t + b_0c_0/a_0)^{\frac{1}{3}}, \\ c(t) = a_0^{\frac{1}{3}} b_0^{-\frac{1}{3}} c_0^{\frac{2}{3}} (\frac{3}{2}t + b_0c_0/a_0)^{\frac{1}{3}}, \end{cases} \tag{3.3}$$

where $a_0 = a(0)$, $b_0 = b(0)$ and $c_0 = c(0)$. In particular, the behavior of this solution is the following:

Lemma 3.1 ([KM01])**.** *The metric (3.1) satisfying (3.2) has three properties.*

1. *The solution of the Ricci flow equation exists on $(-T, +\infty)$, where $T = \frac{b_0c_0}{3a_0}$.*
2. *If $t \to -T$, then $a \to +\infty$, $b \to 0$ and $c \to 0$.*
3. *If $t \to +\infty$, then $a \to 0$, $b \to +\infty$ and $c \to +\infty$.*

We use this lemma later to describe the resulting Ricci-flat metric on the space-time. Next we consider a cohomogeneity one metric with respect to the Heisenberg group H_3. Let (M^4, g) be a cohomogeneity one with respect to H_3. Using the above frame, a cohomogeneity one metric is described as

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{3.4}$$

Put $F_0 = \frac{\partial}{\partial t}$, then $\{F_i\}$ becomes an orthogonal frame. The Ricci tensor of this metric is given by

$$\begin{aligned} R_{00} &= -\frac{\ddot{a}}{a} - \frac{\ddot{b}}{b} - \frac{\ddot{c}}{c}, \\ R_{11} &= -a\frac{(\dot{a}bc)^{\cdot}}{bc} + \frac{a^4}{2b^2c^2}, \\ R_{22} &= -b\frac{(\dot{b}ca)^{\cdot}}{ca} - \frac{a^2}{2c^2}, \\ R_{33} &= -c\frac{(\dot{c}ab)^{\cdot}}{ab} - \frac{a^2}{2b^2}, \end{aligned}$$

and other components are 0. Using (3.3), we construct a Ricci-flat metric in the following way:

Theorem 3.2 ([GR00]). *Let a, b and c satisfy the Ricci flow equations on H_3:*

$$\begin{aligned} \frac{d}{dt}a &= -\frac{a^2}{2bc}, \\ \frac{d}{dt}b &= \frac{a}{2c}, \\ \frac{d}{dt}c &= \frac{a}{2b}. \end{aligned}$$

Then g becomes a Ricci-flat metric.

Proof. An easy computation shows that

$$\begin{aligned} \frac{d^2}{dt^2}a &= \frac{a^3}{b^2c^2}, \\ \frac{d^2}{dt^2}b &= -\frac{a^2}{2bc^2}, \\ \frac{d^2}{dt^2}c &= -\frac{a^2}{2b^2c}, \end{aligned}$$

and ab, ac and b/c are constants. Using this fact and the equation (2.3), we obtain the statement. ☐

Sectional curvatures of this metric are

$$K_{01} = K_{23} = -\frac{a^2}{b^2c^2}, \quad K_{02} = K_{13} = \frac{a^2}{2b^2c^2}, \quad K_{03} = K_{12} = \frac{a^2}{2b^2c^2}.$$

From Proposition (3.1), we get the following proposition.

Proposition 3.3. *Assume that* $a_0b_0c_0$ *is positive. Then*

1. *the parameter* t *exists on* $(-T,\infty)$*, where* $T = \dfrac{2b_0c_0}{3a_0}$.
2. *If* $t \to -T$*, then* $K_{ij} \to \pm\infty$.
3. *If* $t \to +\infty$*, then* $K_{ij} \to 0$.

Assume that $a_0b_0c_0$ *is negative. Then*

1. *the parameter* t *exists on* $(-\infty, T')$*, where* $T' = -\dfrac{2b_0c_0}{3a_0}$.
2. *If* $t \to -\infty$*, then* $K_{ij} \to T'$.
3. *If* $t \to +\infty$*, then* $K_{ij} \to 0$.

Actually, this phenomenon is related with the hyper-Kähler structure. We confirm it from now on. Before we define almost complex structures, we put

$$e^0 = dt, \quad e^1 = a\theta^1, \quad e^2 = b\theta^2, \quad e^3 = c\theta^3.$$

Then $\{e^i\}_{i=0}^3$ becomes an orthonormal coframe. Let $\{e_i\}$ be the orthonormal frame defined by $e^i(e_j) = \delta^i_j$. For $i = 1$, 2 and 3, we consider almost complex structures J_i defined by

$$J_ie_0 = e_i, \quad J_ie_j = -e_k, \quad (J_i)^2 = -id,$$

where (i,j,k) are cyclic permutation of $\{1,2,3\}$. It is easy to check that $\{J_i\}$ satisfy $J_3 = -J_1J_2 = J_2J_1$. Moreover from direct calculation we have $N_{J_i}(F_j, F_k) = 0$ for all $i,j,k = 0,1,2,3$ where N_{J_i} is the Nijenhuis tensor of the almost complex structure J_i. Therefore the almost complex structures $\{J_i\}$ are all integrable. We consider a triple of 2-forms $(\omega^1,\omega^2,\omega^3)$, defined by

$$\omega^i(X,Y) := g(J_iX,Y)$$

for $i = 1,2,3$. Then 2-forms ω_i are given by

$$\omega^i = e^0 \wedge e^i - e^j \wedge e^k,$$

where (i,j,k) are cyclic permutation of $\{1,2,3\}$. The above triple of 2-forms $(\omega^1,\omega^2,\omega^3)$ satisfies the following relations

$$(\omega^i)^2 \neq 0, \quad \omega^i \wedge \omega^j = 0 \quad (i \neq j).$$

Since $de^0 = d(dt) = 0, \quad de^1 = \dot{a}dt \wedge \theta^1 - a\theta^2 \wedge \theta^3, \; de^2 = \dot{b}dt \wedge \theta^2, \quad de^3 = \dot{c}dt \wedge \theta^3$, we get

$$\begin{aligned} d\omega^1 &= (a - \dot{b}c - b\dot{c})dt \wedge \theta^2 \wedge \theta^3, \\ d\omega^2 &= (\dot{a}c + a\dot{c})dt \wedge \theta^3 \wedge \theta^1, \\ d\omega^3 &= (\dot{a}b + a\dot{b})dt \wedge \theta^1 \wedge \theta^2. \end{aligned}$$

Theorem 3.4. *Assume that the cohomogeneity one metric (3.4) has almost complex structures J_i. For any $i = 1, 2, 3$, $d\omega^i = 0$ if and only if a, b and c satisfy the Ricci flow equations on the Heisenberg group.*

Proof. We see at once that $d\omega^i = 0$ if and only if

$$a - \dot{b}c - b\dot{c} = 0, \tag{3.9a}$$

$$\dot{a}c + a\dot{c} = 0, \tag{3.9b}$$

$$\dot{a}b + a\dot{b} = 0. \tag{3.9c}$$

Equations (3.9b) and (3.9c) imply that ac and ab are constants. Using equations (3.9a)-(3.9c), we compute that

$$\begin{aligned} a = (bc)^{\cdot} &= \Big(\frac{ab \cdot ac}{a^2}\Big)^{\cdot} \\ &= \Big(\frac{1}{a^2}\Big)^{\cdot} a^2bc \\ &= -2\dot{a}\frac{bc}{a}. \end{aligned}$$

Hence we get

$$\dot{a} = -\frac{a^2}{2bc}.$$

Since $a\dot{b} = -b\dot{a} = a^2/(2c)$ and $a\dot{c} = -c\dot{a} = a^2/(2b)$, we get

$$\dot{b} = \frac{a}{2c}, \tag{3.11a}$$

$$\dot{c} = \frac{a}{2b}. \tag{3.11b}$$

Therefore a, b and c satisfy the Ricci flow equations on H_3.

Conversely, a, b and c satisfy the Ricci flow equations on H_3, then $d\omega_i = 0$. ☐

Therefore the cohomogeneity one metric with (3.2) becomes the hyper-Kähler metric.

4 The group of rigid motions of the Minkowski 2-space

In this section, we consider the group of rigid motions of the Minkowski 2-space. Let $\{F_i\}_{i=1}^3$ be the Milnor frame of $E(1,1)$, satisfying $[F_1, F_2] = -F_3$, $[F_3, F_1] = 0$, $[F_2, F_3] = F_1$, and $\{\theta_i\}$ the dual coframe of $\{F_i\}$. Then a left-invariant metric is expressed as

$$g = a(\theta^1)^2 + b(\theta^2)^2 + c(\theta^3)^2. \tag{4.1}$$

Therefore the Ricci flow equation is given by

$$\begin{aligned} \frac{d}{dt}a &= \frac{c^2 - a^2}{2bc}, \\ \frac{d}{dt}b &= \frac{(a+c)^2}{2ac}, \\ \frac{d}{dt}c &= \frac{a^2 - c^2}{2ab}. \end{aligned} \tag{4.2}$$

Lemma 4.1 ([KM01])**.** *The Ricci flow on $E(1,1)$ has the following properties.*

1. *The solution of the Ricci flow equation (4.2) exists on $(-T, +\infty)$, where $T > 0$ is depending only on initial data.*
2. *The quantity ac and $b(c-a)$ are conserved.*
3. *Put $\rho := a/c$, then*
$$\begin{aligned} \frac{d}{dt}b &= \frac{(1+\rho)^2}{2\rho}, \\ \frac{d}{dt}\rho &= \frac{1-\rho^2}{b}. \end{aligned}$$
4. $\rho_0 < \rho_\infty \leq 1$ *or* $1 \leq \rho_\infty < \rho_0$.
5. *If $\rho \neq 1$, then*
$$b = k_0 \frac{\sqrt{\rho}}{|1-\rho|},$$
where $k_0 = b_0 \frac{|1-\rho_0|}{\sqrt{\rho_0}}$.

6. *If $t \to +\infty$, then $\rho \to 1$ and $b \to \infty$.*

7. *If $t \to +\infty$, then a and c converge to $a_\infty = c_\infty = \sqrt{a_0 c_0}$.*

We use this proposition later to describe the asymptotic behavior of the resulting Ricci-flat metric on the space-time.

Remark 4.2. If $a = c$, then the metric (4.1) becomes a non-gradient expanding Ricci soliton. See Theorem **??**.

Next, let (M^4, g) be a cohomogeneity one with respect to $E(1,1)$. Let $\{F_i\}_{i=1}^3$ and $\{\theta_i\}$ be as before. Then a cohomogeneity one metric is expressed as

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{4.3}$$

Putting $F_0 = \frac{\partial}{\partial t}$, we have an orthogonal frame $\{F_i\}_{i=1}^4$. The Ricci tensor is computed as

$$\begin{aligned} R_{00} &= -\frac{\ddot{a}}{a} - \frac{\ddot{b}}{b} - \frac{\ddot{c}}{c}, \\ R_{11} &= -a\frac{(\dot{a}bc)^\cdot}{bc} + \frac{a^4 - c^4}{2b^2c^2}, \\ R_{22} &= -b\frac{(\dot{b}ca)^\cdot}{ca} - \frac{(a^2+c^2)^2}{2a^2c^2}, \\ R_{33} &= -c\frac{(\dot{c}ab)^\cdot}{ab} + \frac{c^4 - a^4}{2a^2b^2}, \end{aligned}$$

and other component are 0. The Ricci flow equation for (4.1) is equivalent to the following:

$$\begin{aligned} \frac{d}{dt}a &= \frac{c^2 - a^2}{2bc}, \\ \frac{d}{dt}b &= \frac{(c+a)^2}{2ac}, \\ \frac{d}{dt}c &= \frac{a^2 - c^2}{2ab}. \end{aligned} \tag{4.5}$$

Since $\dfrac{d}{dt}(ac) = 0$, it is easy to check that

$$\begin{aligned} R_{22} &= -b\frac{(\dot{b}ca)^{\cdot}}{ca} - \frac{(c^2+a^2)^2}{2a^2c^2}, \\ &= -b\ddot{b} - \frac{(c^2+a^2)^2}{2a^2c^2}, \\ &= \frac{(c^2-a^2)^2}{2a^2c^2} - \frac{(c^2+a^2)^2}{2a^2c^2} = -2, \end{aligned}$$

and this metric is not a Ricci-flat metric.

Remark 4.3. The metric (4.3) with (4.5) is a Ricci soliton if and only if $a = -c$, in other words, a, b and c are constants. Furthermore, this Ricci soliton is the product metric on $\mathbb{R} \times E(1,1)$ and a non-gradient expanding Ricci soliton. We can check in the following way. First, because $E(1,1) \approx \mathbb{R}^3$, we can use standard coordinates (x_1, x_2, x_3) on $\mathbb{R}^3$. Secondly, since left-invariant vector fields is given by

$$F_1 = e^{x_2}\frac{\partial}{\partial x_3} + e^{-x_2}\frac{\partial}{\partial x_1},\ F_2 = \frac{\partial}{\partial x_2},\ F_3 = e^{x_2}\frac{\partial}{\partial x_3} - e^{-x_2}\frac{\partial}{\partial x_1},$$

then we can write the Ricci soliton equation using standard coordinates (x_1, x_2, x_3) on $\mathbb{R}^3$, and we will obtain conditions of the Ricci soliton, for example,

$$X = \frac{\alpha t}{2}F_0 + e^{-x_2}\frac{\alpha x_3}{4}F_1 + \frac{\alpha}{4}F_2\ ,\ \alpha = \frac{4}{b^2}.$$

Lastly, we check that $\nabla_i X_j$ is not symmetric. Therefore this Ricci soliton is a non-gradient expanding Ricci soliton.

Next, we consider a cohomogeneity one metric

$$g = dt^2 + a(t)^2(\theta^1)^2 - b(t)^2(\theta^2)^2 - c(t)^2(\theta^3)^2. \tag{4.7}$$

This metric has signature $(2,2)$. Put $F_0 = \frac{\partial}{\partial t}$. Then $\{F_i\}$ is the orthogonal frame. The Ricci tensor is given by

$$\begin{aligned}
R_{00} &= -\frac{\ddot{a}}{a} - \frac{\ddot{b}}{b} - \frac{\ddot{c}}{c},\\
R_{11} &= -a\frac{(\dot{a}bc)^{\cdot}}{bc} + \frac{a^4 - c^4}{2b^2c^2},\\
R_{22} &= b\frac{(\dot{b}ca)^{\cdot}}{ca} + \frac{(c^2 - a^2)^2}{2a^2c^2},\\
R_{33} &= c\frac{(\dot{c}ab)^{\cdot}}{ab} + \frac{a^4 - c^4}{2a^2b^2}.
\end{aligned}$$

These formulas imply the following theorem:

Theorem 4.4. *Let a, b and c satisfy the Ricci flow equations;*

$$\begin{aligned}
\frac{d}{dt}a &= \frac{c^2 - a^2}{2bc},\\
\frac{d}{dt}b &= \frac{(c + a)^2}{2ca},\\
\frac{d}{dt}c &= \frac{a^2 - c^2}{2ab}.
\end{aligned} \tag{4.9}$$

Then the metric (4.7) becomes a Ricci-flat metric.

Proof.

$$\begin{aligned}
\frac{d^2}{dt^2}a &= \frac{(a^2 - c^2)(2a^2 + ac + c^2)}{ab^2c^2},\\
\frac{d^2}{dt^2}b &= -\frac{(c^2 - a^2)^2}{2a^2bc^2},\\
\frac{d^2}{dt^2}c &= \frac{(a^2 - c^2)(a^2 + ac + 2c^2)}{2a^2b^2c},
\end{aligned} \tag{4.10}$$

and ac is a constant. Using this fact, we obtain the statement. □

Sectional curvatures of the metric (4.7) satisfying (4.2) are computed as

$$\begin{aligned}
K_{01} = K_{23} &= \frac{(c^2 - a^2)(2a^2 + ac + c^2)}{2a^2b^2c^2} = \frac{(1 - \rho^2)(2\rho^2 + \rho + 1)}{2\rho^2b^2},\\
K_{02} = K_{13} &= \frac{(c^2 - a^2)^2}{2a^2b^2c^2} = \frac{(1 - \rho^2)^2}{2\rho^2b^2},\\
K_{03} = K_{12} &= -\frac{(c^2 - a^2)(a^2 + ac + 2c^2)}{2a^2b^2c^2} = -\frac{(1 - \rho^2)(\rho^2 + \rho + 2)}{2\rho^2b^2},
\end{aligned}$$

where $\rho := a/c$. Notice that if $a^2 \neq c^2$, then signatures of sectional curvatures K_{ij} are decided. From Proposition (4.1), we get the following proposition.

Proposition 4.5. *Let a, b and c satisfy the Ricci flow equations (4.2).*

1. *If $a^2 = c^2$, then the metric (4.7) becomes a zero curvature metric.*
2. *If $a^2 \neq c^2$, then the metric (4.7) is a Ricci-flat metric, but K_{ij} are not constant.*

Assume that $a_0 c_0$ and b_0 are positive.

1. *The Ricci flow exists on $(-T, \infty)$, where T depends on the initial data.*
2. *If $t \to +\infty$, then all sectional curvatures $K_{ij} \to 0$.*
3. *If $t \to -T$, then $K_{ij} \to \pm\infty$.*

Assume that $a_0 c_0$ is positive, and b_0 is negative.

1. *The Ricci flow exists on $(-\infty, T)$, where T depends on initial data.*
2. *If $t \to -T$, then sectional curvature $K_{ij} \to 0$.*
3. *If $t \to \infty$, then $K_{ij} \to \pm\infty$.*

Assume that $a_0 c_0$ is negative, and b_0 is positive. Then the behavior of a, b and c is the backward Ricci flow of $E(2)$ (see (5.1)).

Assume that $a_0 c_0$ and b_0 are negative. Then the behavior of a, b and c is the Ricci flow of $E(2)$ (see (5.1)).

5 The group of rigid motions of the Euclidean 2-space

In this section, we consider the group $E(2)$ of rigid motions of the Euclidean 2-space. Let $\{F_i\}_{i=1}^3$ be the Milnor frame of $E(2)$, satisfying

$$[F_2, F_3] = -F_1 \ , \ [F_3, F_1] = 0 \ , \ [F_1, F_2] = -F_3,$$

and $\{\theta_i\}$ dual coframe of $\{F_i\}$. Then a left-invariant metric is expressed as

$$g = a(\theta^1)^2 + b(\theta^2)^2 + c(\theta^3)^2.$$

Therefore the Ricci flow equation is equivalent to

$$\begin{cases} \dfrac{d}{dt}a = \dfrac{c^2 - a^2}{2bc}, \\ \dfrac{d}{dt}b = \dfrac{(c-a)^2}{2ca}, \\ \dfrac{d}{dt}c = \dfrac{a^2 - c^2}{2ab}. \end{cases} \tag{5.1}$$

Since $E(1,1)$ and $E(2)$ are solvable, so the Ricci flow equations (5.1) resemble as the Ricci flow equations (4.2). The behavior of the solution of (5.1) is known in [KM01, CK04].

Lemma 5.1 ([KM01, CK04])**.** *The Ricci flow on $E(2)$ has the following properties.*

1. *The solution of the the Ricci flow equation (5.1) exists on $(-T, \infty)$, where $T > 0$ is depending only on initial data.*
2. *The quantities ac and $b(c + a)$ are preserved.*
3. *Put $k := a/c$. Then we get*

$$\frac{d}{dt}b = \frac{(1-k)^2}{2k},$$

$$\frac{d}{dt}k = \frac{1-k^2}{b}.$$

4. *If $t \to \infty$, then $k \to 1$ and $b \to^{\exists} b_\infty$.*
5. *The coefficient b satisfies*

$$b = l_0 \frac{\sqrt{k}}{1+k},$$

where $l_0 = b_0 \frac{1+k_0}{\sqrt{k_0}}$.

We use above proposition later to describe the asymtptotic property of the resulting Ricci-flat metric on the space-time.

Remark 5.2. Since $E(2)$ is dynamically stable, the Ricci flow on $E(2)$ converges to an Einstein metric. The details are written in [IJ92, Ses06, SSS08].

Next, we consider a cohomogeneity one metric with respect to $E(2)$. Let $\{F_i\}_{i=1}^3$ and $\{\theta_i\}$ be as before. Then the metric is expressed as

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{5.2}$$

The Ricci tensor of this metric is given by

$$\begin{aligned}
R_{00} &= -\frac{\ddot{a}}{a} - \frac{\ddot{b}}{b} - \frac{\ddot{c}}{c},\\
R_{11} &= -a\frac{(\dot{a}bc)^{\cdot}}{bc} + \frac{a^4 - c^4}{2b^2c^2},\\
R_{22} &= -b\frac{(\dot{b}ca)^{\cdot}}{ca} - \frac{(a^2 - c^2)^2}{2a^2c^2},\\
R_{33} &= -c\frac{(\dot{c}ab)^{\cdot}}{ab} + \frac{c^4 - a^4}{2a^2b^2},
\end{aligned}$$

and other components are 0. Using the solution of (5.1), we construct a Ricci-flat metric on the space-time in the following way:

Theorem 5.3. *If a, b and c satisfy the Ricci flow equations*

$$\begin{cases}
\dfrac{d}{dt}a = \dfrac{c^2 - a^2}{2bc},\\
\dfrac{d}{dt}b = \dfrac{(a - c)^2}{2ac},\\
\dfrac{d}{dt}c = \dfrac{a^2 - c^2}{2ab},
\end{cases} \tag{5.4}$$

then this metric (5.2) becomes a Ricci-flat metric.

Proof. This is proved by routine calculations. ☐

Sectional curvatures of this metric are

$$\begin{aligned}
K_{01} = K_{23} &= \frac{(c^2 - a^2)(2a^2 - ac + c^2)}{a^2b^2c^2} = \frac{(1 - k^2)(2k^2 - k + 1)}{2b^2k^2},\\
K_{02} = K_{13} &= \frac{(c^2 - a^2)^2}{2a^2b^2c^2} = \frac{(1 - k^2)^2}{2b^2k^2},\\
K_{03} = K_{12} &= -\frac{(c^2 - a^2)(a^2 - ca + 2c^2)}{2a^2b^2c^2} = -\frac{(1 - k^2)(k^2 - k + 2)}{2b^2k^2},
\end{aligned}$$

where $k := a/c$. We can observe similarities between the Ricci flow solutions on the space-times of the left-invariant Ricci flow solutions of $E(1, 1)$ and $E(2)$. For instance, we get the following proposition.

Proposition 5.4.

1. *If $a^2 = c^2$, then this metric is a flat curvature metric.*
2. *If $a^2 \neq c^2$, then this metric is a Ricci-flat metric but not a constant curvature metric.*

Let a_0c_0 and b_0 be positive.

1. *The Ricci flow equation exists on $(-T, \infty)$, where T depends on initial data.*
2. *If $t \to +\infty$, then $K_{ij} \to 0$.*

Let a_0c_0 be positive, and b_0 be negative.

1. *The Ricci flow equation exists on $(-\infty, T)$, where T depends on initial data.*
2. *If $t \to -T$, then sectional curvatures $K_{ij} \to 0$.*
3. *If $t \to \infty$, then $K_{ij} \to \pm\infty$.*

Let a_0c_0 be negative, and b_0 be positive. Then behavior of a, b and c is the backward Ricci flow of $E(1,1)$ (see (4.2)). Let a_0c_0 and b_0 be negative. Then behavior of a, b and c is the Ricci flow of $E(1,1)$ (see (4.2)).

Next, we define almost complex structures. Using $E(2) \approx \mathbb{R}^3$, we can write

$$\begin{aligned}
F_1 &= \sin y \frac{\partial}{\partial x} + \cos y \frac{\partial}{\partial z}, \\
F_2 &= \frac{\partial}{\partial y}, \\
F_3 &= \cos y \frac{\partial}{\partial x} - \sin y \frac{\partial}{\partial z},
\end{aligned}$$

and

$$\begin{aligned}
\theta^1 &= \sin y \cdot dx + \cos y \cdot dz, \\
\theta^2 &= dy, \\
\theta^3 &= \cos y \cdot dx - \sin y \cdot dz.
\end{aligned}$$

Then the metric on $E(2)$ with $a = b = c = 1$ is

$$g_3 = (\theta^1)^2 + (\theta^2)^2 + (\theta^3)^2 = dx^2 + dy^2 + dz^2.$$

Let $\{e_i\}_{i=0}^3$ be an orthonormal coframe defined by

$$e^0 = dt\ ,\ e^1 = a\theta^1\ ,\ e^2 = b\theta^2\ ,\ e^3 = c\theta^3.$$

Then $\{e^i\}_{i=0}^4$ becomes an orthonormal coframe of cohomogeneity one metric g. Let $\{e_i\}$ be the orthonormal frame, satisfying $e^i(e_j) = \delta^i_j$. Since $\mathbb{R}^4 \approx \mathbb{C}^2$ has natural almost complex structures $\{J_i\}$, we analogously define almost complex structures J_1 and J_2, satisfying

$$\begin{aligned} J_1 e_0 &= e_2, \quad J_1 e_3 = e_1, \quad (J_1)^2 = -id, \\ J_2 e_0 &= \cos y \cdot e_1 - \sin y \cdot e_3, \\ J_2 e_2 &= \sin y \cdot e_1 + \cos y \cdot e_3, \\ (J_2)^2 &= -id. \end{aligned}$$

If we put $J_3 := J_2 J_1 = -J_1 J_2$, then J_3 becomes an almost complex structure. From direct computation we have $N_{J_i}(F_j, F_k) = 0$ for all $i, j, k = 0, 1, 2, 3$ where N_{J_i} is the Nijenhuis tensor of J_i. Therefore J_i's are all integrable. We consider a triple of 2-forms $(\omega^1, \omega^2, \omega^3)$, defined by

$$\omega^i(X, Y) := g(J_i X, Y)$$

for $i = 1, 2, 3$. Then 2-forms ω_i are given by

$$\begin{aligned} \omega^1 &= e^0 \wedge e^2 + e^3 \wedge e^1, \\ \omega^2 &= \cos y \cdot (e^0 \wedge e^1 + e^2 \wedge e^3) - \sin y \cdot (e^0 \wedge e^3 + e^1 \wedge e^2), \\ \omega^3 &= \sin y \cdot (e^0 \wedge e^1 + e^2 \wedge e^3) + \cos y \cdot (e^0 \wedge e^3 + e^1 \wedge e^2). \end{aligned}$$

The above triple of 2-forms $(\omega^1, \omega^2, \omega^3)$ satisfies the following relations:

$$(\omega^i)^2 \neq 0,\ \omega^i \wedge \omega^j = 0\ (i \neq j).$$

Since $de^0 = d(dt) = 0,\ de^1 = \dot{a} dt \wedge \theta^1 + a\theta^2 \wedge \theta^3,\ de^2 = \dot{b} dt \wedge \theta^2,\ de^3 = \dot{c} dt \wedge \theta^3 + c\theta^1 \wedge \theta^2$, we get

$$\begin{aligned} d\omega^1 &= (\dot{c}a + c\dot{a}) dt \wedge \theta^3 \wedge \theta^1, \\ d\omega^2 &= -(\dot{a}b + a\dot{b} + a - c) \sin y \cdot dt \wedge \theta^1 \wedge \theta^2 + (\dot{b}c + b\dot{c} + c - a) \cos y \cdot dt \wedge \theta^2 \wedge \theta^3, \\ d\omega^3 &= -(\dot{a}b + a\dot{b} + a - c) \cos y \cdot dt \wedge \theta^1 \wedge \theta^2 + (\dot{b}c + b\dot{c} + c - a) \sin y \cdot dt \wedge \theta^2 \wedge \theta^3. \end{aligned}$$

Theorem 5.5. *The cohomogeneity one metric g expressed as (5.2), where the triple $\{a(t), b(t), c(t)\}$ satisfies the Ricci flow equation on $E(2)$, is a hyper-Kähler metric. Conversely the cohomogeneity one metric g as in (5.2) satisfies $d\omega^i = 0$ for any $i = 1, 2, 3$, then the triple $\{a(t), b(t), c(t)\}$ satisfies the Ricci flow equations on $E(2)$.*

Proof. It is straitforward to check that $d\omega^i = 0$ if and only if the following hold:

$$\dot{c}a + c\dot{a} = 0, \tag{5.11a}$$

$$\dot{a}b + a\dot{b} = c - a, \tag{5.11b}$$

$$\dot{b}c + b\dot{c} = a - c. \tag{5.11c}$$

(5.11a) implies that ca is constant. Solveing the equations for $(\dot{a}, \dot{b}, \dot{c})$, we obtain the statement of the theorem. ☐

Therefore the cohomogeneity one metric (5.4) becomes a hyper-Kähler metric.

Remark 5.6. In this section, we obtain the cohomogeneity one metric with respect to $E(2)$ with the hyper-Kähler structure $\{J_i\}$. From the definition of $\{J_i\}$, the hyper-Kähler structure is non-trivial. The cohomogeneity one metric with respect to $E(2)$ with the trivial hyper-Kähler structure (i.e. $J_ie_0 = e_i, J_ie_j = e_k$, where (i, j, k) are cyclic permutation of $\{1, 2, 3\}$,) is also a hyper-Kähler structure, however the triple of functions $\{a(t), b(t), c(t)\}$ does not satisfy the Ricci flow equation. So we do not consider it deeply.

6 $SL(2, \mathbb{R})$

In this section, we consider $SL(2, \mathbb{R})$. Let (M^4, g) be a cohomogeneity one with respect to $SL(2, \mathbb{R})$. Let $\{F_i\}_{i=1}^3$ be the Milnor frame of $SL(2, \mathbb{R})$, satisfying

$$[F_1, F_2] = -F_3, \ [F_2, F_3] = F_1, \ [F_3, F_1] = F_2,$$

and $\{\theta_i\}$ the dual coframe of $\{F_i\}$. The metric is expressed as

$$g = dt^2 + a(t)^2(\theta^1)^2 + b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{6.1}$$

The Ricci tensor of this metric is given by

$$\begin{aligned}
R_{00} &= -\frac{\ddot{a}}{a} - \frac{\ddot{b}}{b} - \frac{\ddot{c}}{c},\\
R_{11} &= -a\frac{(\dot{a}bc)^{\cdot}}{bc} - \frac{(b^2+c^2)^2-a^4}{2b^2c^2},\\
R_{22} &= -b\frac{(\dot{b}ca)^{\cdot}}{ca} - \frac{(a^2+c^2)^2-b^4}{2a^2c^2},\\
R_{33} &= -c\frac{(\dot{c}ab)^{\cdot}}{ab} - \frac{(a^2-b^2)^2-c^4}{2a^2b^2},
\end{aligned}$$

and other component are 0.

Proposition 6.1. *If a, b and c satisfy the Ricci flow equations of $SL(2,\mathbb{R})$*

$$\begin{cases}
\dfrac{d}{dt}a = \dfrac{(b+c)^2-a^2}{2bc},\\
\dfrac{d}{dt}b = \dfrac{(c+a)^2-b^2}{2ca},\\
\dfrac{d}{dt}c = \dfrac{(a-b)^2-c^2}{2ab},
\end{cases} \tag{6.3}$$

then this metric (6.1) satisfying (6.3) is not a Ricci-flat metric.

Proof. The proof is a straightforward calculation. □

Next, we consider the metric of the form

$$g = dt^2 - a(t)^2(\theta^1)^2 - b(t)^2(\theta^2)^2 + c(t)^2(\theta^3)^2. \tag{6.4}$$

This metric (6.4) has signature $(2,2)$.

The Ricci tensor of this metric is computed as

$$\begin{aligned}
R_{00} &= -\frac{\ddot{a}}{a} - \frac{\ddot{b}}{b} - \frac{\ddot{c}}{c},\\
R_{11} &= a\frac{(\dot{a}bc)^{\cdot}}{bc} + \frac{(b^2-c^2)^2-a^4}{2b^2c^2},\\
R_{22} &= b\frac{(\dot{b}ca)^{\cdot}}{ca} + \frac{(a^2-c^2)^2-b^4}{2a^2c^2},\\
R_{33} &= -c\frac{(\dot{c}ab)^{\cdot}}{ab} - \frac{(a^2-b^2)^2-c^4}{2a^2b^2},
\end{aligned}$$

and other components are 0. The Ricci flow equations of $SL(2,\mathbb{R})$ are given by

$$\begin{aligned}\frac{d}{dt}a &= \frac{(b+c)^2-a^2}{2bc},\\ \frac{d}{dt}b &= \frac{(c+a)^2-b^2}{2ca},\\ \frac{d}{dt}c &= \frac{(a-b)^2-c^2}{2ab}.\end{aligned} \tag{6.6}$$

In this case, changing a into $-a$ and b into $-b$, therefore the Ricci flow equations of $SL(2,\mathbb{R})$ change into

$$\begin{aligned}\frac{d}{dt}a &= \frac{(b-c)^2-a^2}{2bc},\\ \frac{d}{dt}b &= \frac{(c-a)^2-b^2}{2ca},\\ \frac{d}{dt}c &= \frac{(a-b)^2-c^2}{2ab}.\end{aligned} \tag{6.7}$$

This system of equations is nothing but the Ricci flow equations of left-invariant metrics on $SU(2)$.

Theorem 6.2. *If a, b and c satisfy the Ricci flow equations of $SU(2)$, then the metric (6.4) satisfying (6.7) becomes a Ricci-flat metric of signature $(2,2)$.*

Proof. The proof method of this theorem is similar to Theorem 1.3. The proof is straightforward. □

References

[Bes08] A. L. Besse, *Einstein manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition.

[BD07] P. Baird and L. Danielo, *Three-dimensional Ricci solitons which project to surfaces*, J. Reine Angew. Math. **608** (2007), 65–91.

[CGLP04a] M. Cvetič, G. W. Gibbons, H. Lü, and C. N. Pope, *New cohomogeneity one metrics with Spin(7) holonomy*, J. Geom. Phys. **49** (2004), no. 3-4, 350–365.

[CGLP04b] ______, *Orientifolds and slumps in G_2 and* Spin(7) *metrics*, Ann. Physics **310** (2004), no. 2, 265–301.

[CK04] B. Chow and D. Knopf, *The Ricci flow: an introduction*, Mathematical Surveys and Monographs, vol. 110, American Mathematical Society, Providence, RI, 2004.

[DVV92] R. Dijkgraaf, H. Verlinde, and E. Verlinde, *String propagation in a black hole geometry*, Nuclear Phys. B **371** (1992), no. 1-2, 269–314.

[EGH80] T. Eguchi, P. B. Gilkey, and A. J. Hanson, *Gravitation, gauge theories and differential geometry*, Phys. Rep. **66** (1980), no. 6, 213–393.

[Fri85] D. H. Friedan, *Nonlinear models in* $2+\varepsilon$ *dimensions*, Ann. Physics **163** (1985), no. 2, 318–419.

[GIK06] C. Guenther, J. Isenberg, and D. Knopf, *Linear stability of homogeneous Ricci solitons*, Int. Math. Res. Not. (2006), Art. ID 96253, 30.

[GR00] G. W. Gibbons and P. Rychenkova, *Single-sided domain walls in M-theory*, J. Geom. Phys. **32** (2000), no. 3, 311–340.

[Ham82] R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. **17** (1982), no. 2, 255–306.

[IJ92] J. Isenberg and M. Jackson, *Ricci flow of locally homogeneous geometries on closed manifolds*, J. Differential Geom. **35** (1992), no. 3, 723–741.

[Kro89] P. B. Kronheimer, *The construction of ALE spaces as hyper-Kähler quotients*, J. Differential Geom. **29** (1989), no. 3, 665–683.

[Koi90] N. Koiso, *On rotationally symmetric Hamilton's equation for Kähler-Einstein metrics*, Recent topics in differential and analytic geometry, Adv. Stud. Pure Math., vol. 18, Academic Press, Boston, MA, 1990, pp. 327–337.

[KM01] D. Knopf and K. McLeod, *Quasi-convergence of model geometries under the Ricci flow*, Comm. Anal. Geom. **9** (2001), no. 4, 879–919.

[Lor83] D. Lorenz, *Gravitational instanton solutions for Bianchi types* $\mathrm{I}--\mathrm{IX}$, Acta Phys. Polon. B **14** (1983), no. 11, 791–805.

[LP84] D. Lorenz-Petzold, *On the general vacuum and stiff matter solutions for "diagonal" Bianchi type-*VI_0 *and type-*VII_0 *models*, Acta Phys. Polon. B **15** (1984), no. 2, 117–122. MR746487 (85h:83020)

[LP89] D. Lorenz-Petzold, *Gravitational instanton solutions*, Progr. Theoret. Phys. **81** (1989), no. 1, 17–22.

[Lau01] J. Lauret, *Ricci soliton homogeneous nilmanifolds*, Math. Ann. **319** (2001), no. 4, 715–733.

[Lot07] J. Lott, *On the long-time behavior of type-III Ricci flow solutions*, Math. Ann. **339** (2007), no. 3, 627–666.

[Mil76] J. Milnor, *Curvatures of left invariant metrics on Lie groups*, Advances in Math. **21** (1976), no. 3, 293–329.

[Nom81] K. Nomizu, *Introduction to Modern Differential Geometry*, Interscience Tracts in Pure and Applied Mathematics, No. 15 Vol. II, Shoka-bo, 1981 (Japaneses).

[Nod08] T. Noda, *A special Lagrangian fibration in the Taub-NUT space*, J. Math. Soc. Japan **60** (2008), no. 3, 653–663.

[Per02] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, preprint: math. DG/0211159 (2002).

[Ses06] N. Sesum, *Linear and dynamical stability of Ricci-flat metrics*, Duke Math. J. **133** (2006), no. 1, 1–26.

[SSS08] O. C. Schnürer, F. Schulze, and M. Simon, *Stability of Euclidean space under Ricci flow*, Comm. Anal. Geom. **16** (2008), no. 1, 127–158.

[TY90] G. Tian and S.-T. Yau, *Complete Kähler manifolds with zero Ricci curvature. I*, J. Amer. Math. Soc. **3** (1990), no. 3, 579–609.

[Ura] H. Urakawa, *Equivariant theory of Einstein metrics on Riemannian manifolds of cohomogeneity one*, Unpublished.